

La Asociación de Geólogos y Geofísicos Españoles del Petróleo (AGGEP)

Os invita a la conferencia que con el título:

“Sequence Stratigraphy and Facies Architecture of Deep-water Reservoirs: New Data, Tools and Concepts”




Impartirá

Timothy R. Garfield

Distinguished Lecturer de la AAPG

Y tendrá lugar en el Auditorio de Repsol-Ypf ,
Pº de la Castellana, 278, el día:

Martes, 1 de Abril de 2003, a las 18:00 horas



2002-03 AAPG Distinguished Lecture

Abstract

TIMOTHY R. GARFIELD

ExxonMobil Exploration Co.
Houston, TX

Funded by the AAPG Foundation Dean A. McGee endowment

Sequence Stratigraphy and Facies Architecture of Deep-water Reservoirs: New Data, Tools and Concepts

Current understanding of deep-water siliciclastics arrived relatively late on the stage of sedimentary geology. Sedimentary processes active at bathyal to abyssal depths did not lend themselves to empirical analog study so important to the early advancements achieved in other depositional settings. As a result, understanding the stratigraphy and facies architecture of deep-water reservoirs has been historically linked to and dependent on technology development. Because direct observations of deep-ocean turbidity currents and debris flows was, until recently, impossible; conceptual understanding of how these sediments were transported and deposited had to await the development of new remote sensing tools and technologies. For example, the first indirect indications of large scale, naturally occurring sediment gravity flows followed the development and distribution of sea-floor communications cables in the late 1800's and early 1900's. Periodic cable breakages demonstrated the existence of turbidity currents and allowed early researchers to estimate their power and velocity. Later, the development of experimental flumes lead to better understanding of the flow mechanisms associated with graded beds common to "Flycsh" deposits, now known to be dominantly turbidites.

In the 1970's and early '80's, several nearly identical models for submarine fans were developed, based largely on outcrop studies. These models described an inner (or upper) channelized fan, a lobe dominated mid fan and an extensive thin-bedded, fine-grained sheet deposit comprising the outer fan. Submarine fan models have evolved dramatically with the advent of high-resolution 3-D seismic, side-scan sonar and other water-bottom imaging tools developed in the 1980's and 90's. Extensive channel networks are now known to be persistent in deep-water systems and channelized flow more prevalent and important to the development of reservoir-grade sand deposits relative to unconfined flow. Channels play an important role in moving sands to even the most distal reaches of many submarine fans, whereas low density silts and muds are commonly stripped off the

upper part of turbidity currents and deposited as overbank in more proximal to medial areas dominated by slope channels. This flow stripping process often results in an overall proximal to distal increase in the sand/mud ratio, the inverse of sediment distribution patterns depicted in the classic fan models.

The development of seismic stratigraphy and later sequence stratigraphy concepts in the 1970's and 1980's was based largely on observations made on multi-channel 2D seismic, well and outcrop data from passive margin settings around the world. Based on principals of accommodation, sequence stratigraphy has provided a conceptual framework to better understand and predict the stratigraphic occurrence, distribution and facies variability of deep-water reservoirs. Improved imaging using modern 3D seismic and the deep-water drilling activities of the 1990's has confirmed the fundamental principals of sequence stratigraphy and allowed the concepts to be applied to deep-water reservoir interpretation and prediction at increasingly higher levels of resolution. These data show that a lowstand cycle of sediment flux may generate early slope failure, slump and slide deposits associated with falling sea level. Early slope failure deposits are often overlain by coarser-grained distributary channel sands, which are primary reservoir targets. These sand-rich units are mostly deposited during peak lowstand times when sediment flux and shelf by-pass are at a maximum. Late-stage deposition of progressively finer-grained, levee-associated turbidites is a common product of waning coarse sediment supply to the upper slope as shelf accommodation increases during early transgression. This idealized sequence stacking pattern is predictive, but not always observed at every location along the depositional profile-because sites of deposition tend to shift both laterally and first basinward than landward through time. In addition, a complete depositional stacking pattern is more typical where the grain size input to the system is broadly distributed, i.e. mixed sand/mud systems. Coarse-grained sand-dominated systems typically lack well-developed levees whereas mud

American Association of Petroleum Geologists

An International Geological Organization

P.O. Box 979 • Tulsa, Oklahoma 74101-0979 USA

(918) 560-2621 • Fax; (918) 560-2678 • email: Cammy@aapg.org

dominated systems display well expressed levees and less voluminous distributary elements.

Many of the recent advancements in the data, tools and concepts applied to deep-water siliciclastics have been driven by the need to explore for and develop hydrocarbon reservoirs in the high-cost deep-water environment. For example, recent improvements in 3D-visualization technology have significantly enhanced our understanding of deep-water channel architecture. Previously unimagined, the complex vertical and lateral facies variations associated with temporally evolving and laterally migrating sinuous deep-water channel systems

can now be imaged in great detail. This detailed understanding of channel architecture enables improved predictions of reservoir facies and net:gross, more accurate placement of reservoir properties in to 3D geologic models and more precise positioning of exploration and development wells. With its vast technological resources, extensive global databases and exploration and development focus on deep-water reservoirs, the petroleum industry, in effective partnership with academic research, has contributed greatly to the improved understanding of the stratigraphy and facies architecture of deep-water reservoirs we now enjoy.

Timothy R. Garfield

EDUCATION:

University of Colorado, Boulder; B.A. in Geology
University of Colorado, Boulder; M.S. in Geology

EXPERIENCE:

1981- 1982 Continental Laboratories, Oklahoma City, Oklahoma; Well-site geologist
1983-1984 Gallagher R&D Co., Lakewood, Colorado; Staff Geologist
1984-1990 RPI International Inc., Boulder, Colorado; Research Geologist, Project Manager
1990- 1995 Exxon Production Research Co., Houston, Texas; Research Geoscientist
1995- 1998 Exxon Production Research Co., Houston, Texas; Group Leader - Deep-Water Reservoirs Group
1998- 2000 Exxon Exploration Co., Houston, Texas; Supervisor, Critical Technology Integrated Reservoir Interpretation Core Group
2000- present ExxonMobil Exploration Co., Houston, Texas; Upstream Geoscience, Stratigraphy Advisor

PUBLICATIONS:

Garfield, T. R., A. J. Scott, and A. L. Walker, 1988, Overview; late Paleozoic geologic history of the Northern Denver Basin, *in* S.M. Goolsby and M. L. Longman, eds., Occurrence and Petrophysical Properties of Carbonate Reservoirs in the Rocky Mountain Region: Rocky Mountain Association of Geologists Publication, p. 1-18.
Garfield, T. R., and M. W. Longman, 1989, Depositional variations in the Fusselman Formation, central Midland Basin, West Texas, *in* B. K. Cunningham and D. W. Cromwell, eds., The Lower Paleozoic of West Texas and Southern New Mexico: Modern Exploration Concepts, Society of Economic Paleontologists and Mineralogists, Permian Basin Chapter Publication, p. 187-202.
Garfield, T. R., N. F. Hurley, and D. A. Budd, 1989, Characterization of pore geometry and natural fractures, Little Sand Draw Field, Big Horn Basin, Wyoming: Abstract, AAPG Bulletin, v. 73, p. 1156.
Garfield, T. R., 1990, Reservoir geology of the upper Phosphoria Formation (Permian), Little Sand Draw

Field, Big Horn Basin, Wyoming: Master's thesis, University of Colorado, Boulder, Colorado, 199 p.
Garfield, T. R., N. F. Hurley, and D. A. Budd, 1992, Little Sand Draw Field, Big Horn Basin, Wyoming: a hybrid dual-porosity and single-porosity reservoir in the Phosphoria Formation: AAPG Bulletin, v. 76 (3), p. 317-391.
Garfield, T. R., D. C. Jennette, F. J. Goulding, and D. K. Sickafoose, 1998, A successful approach to deep-water reservoir prediction; integration of outcrop, 3-D seismic, well-log and core data, and process-response modeling: Abstract, AAPG Bulletin, v. 82, p. 1916-1917.
Garfield, T. R., D. C. Jennette, F. J. Goulding, and D. K. Sickafoose, 1999, An integrated approach to deep-water reservoir prediction: Abstract, AAPG Bulletin, v. 83, p. 1314, 1999.
Garfield, T. R., R. T. Beaubouef, D. C. Jennette, D. C. Mohrig, and A. R. Sprague, 2000, New insight into the three-dimensional architecture of deep-water facies: The product of a multidisciplinary approach, *in* C. Appi, R.S. F. d'Avila and A. R. Viana, eds., Deep-Water Sedimentation: Challenges for the Next Millennium: 31st International Geological Congress Workshop Proceedings, Addendum p. 1-9.
Garfield, T. R., R. T. Beaubouef, and D. K. Sickafoose, 2000, High-resolution Sequence Stratigraphy: A tool for improved reservoir delineation in passive margin settings — Examples from the divergent margin of the South Atlantic Salt Basin: 31st International Geological Congress, Abstracts volume - CD.
Hurley, N. F., T. R. Garfield, S. P. Simmons, 1990, Integrated analysis of natural fractures in Big Horn Basin anticlines: Abstract, AAPG Bulletin, v. 74, p. 680.
Clark, W. J., P. W. Choquette, E. E. Hiatt, T. R. Garfield, D. A. Budd, N. F. Hurley, 1991, Sequence stratigraphy interpretation of the Ervay Member (Permian, Park City Formation), west-central Wyoming paleoshelf : Abstract, AAPG Bulletin, v. 75 , p. 553.
Donovan, A. D., A. W. Djacic, N. S. Ioannides, T. R. Garfield, C. R. Jones, 1992, Sequence stratigraphic control on Middle and Upper Jurassic reservoir distribution within the UK central North Sea, *in* J. P. Parker, J. R., ed., Petroleum Geology of Northwest Europe: Proceedings of the 4th Barbican Conference, p. 251-269.

Beaubouef, R. T., T. R. Garfield, and F. J. Goulding, 1998, Seismic stratigraphy of depositional sequences; high resolution images from a passive margin slope setting, offshore West Africa: Abstract: AAPG Bulletin, v 82, p. 1890

Jennette, D. C., T. R. Garfield, G. W. Farquharson, G. T. Cayley, Glen T., 1998, Facies, architecture and sequence stratigraphy of sand-rich basin floor fans: Examples from the Paleogene, UK central North Sea Anonymous, Abstracts, AAPG Bulletin, v. 82, p. 1926-1927.

Sykes, M. A., C. Rossen, D. C. Mohrig, and T. R. Garfield, 1998 Lithofacies associations within complex slope channel reservoirs; debrites and turbidites: Abstract, AAPG Bulletin, v. 82, p. 1973.

Friedmann, S. J., T. R. Garfield, and D. C. Jennette, 1999 Morphology of channels, fans, and intraslope basins, Late Pleistocene of the West Niger Delta; implications for sedimentological processes and what drives them: AAPG Annual Meeting, Expanded Abstracts, p. A44

Jennette, D. C. and T. R. Garfield, 1999, The interaction of shelf accommodation, sediment supply and sea level in controlling the facies, architecture and distribution of

the Forties and Tay basin-floor fans, central North Sea: Abstract, AAPG Bulletin, v. 83, p. 1320.

Goulding, F. J., T. R. Garfield, K. W. Rudolph, G. N. Jensen, and R. T. Beaubouef, 2000, Seismic/sequence stratigraphy of deep-water reservoirs; 1, Seismic facies and recognition criteria; Past experiences and new observations: AAPG Annual Meeting, Expanded Abstracts, p. 56.

MEMBERSHIPS:

American Association of Petroleum Geologists

RESEARCH INTERESTS:

Seismic stratigraphy and facies analysis of deepwater reservoirs. Lead several regional exploration projects focused on deepwater reservoirs of the North Sea, offshore Nigeria and Angola. Results from these projects contributed to improve sequence stratigraphy and facies models for deepwater systems. Currently contributes to deepwater reservoir evaluations for ExxonMobil's deepwater exploration, development and production projects worldwide.